

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷: A61K 31/505, A61P 25/16	A1	(11) International Publication Number: WO 00/16777 (43) International Publication Date: 30 March 2000 (30.03.00)
(21) International Application Number: PCT/IB99/01503 (22) International Filing Date: 3 September 1999 (03.09.99) (30) Priority Data: 60/101,191 21 September 1998 (21.09.98) US 60/103,868 12 October 1998 (12.10.98) US (71) Applicant (for all designated States except US): PFIZER PRODUCTS INC. [US/US]; Eastern Point Road, Groton, CT 06340 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): JACKSON, Elise, Rose [US/US]; 36 Case Street, Norwich, CT 06360 (US). MCLEAN, Stafford [US/US]; 368 Mistuxet Avenue, Ston- ington, CT 06378 (US). ZORN, Stevin, Howard [US/US]; 62C Patricia Avenue, North Stonington, CT 06359 (US). (74) Agents: SPIEGEL, Allen, J. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: PHARMACEUTICAL AGENTS FOR THE TREATMENT OF PARKINSON'S DISEASE, ADHD AND MICROADENOMAS		
(57) Abstract The present invention is directed to the use of certain pyrido[1,2-a]-pyrazine derivatives, also described as bis-azabicyclic compounds and defined by the formula (I) herein, in the treatment of Parkinson's disease, attention deficit hyperactivity disorder and microadenomas in mammals, and to related compositions.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

5

**PHARMACEUTICAL AGENTS FOR THE TREATMENT
OF PARKINSON'S DISEASE, ADHD AND MICROADENOMAS**

Field of the Invention

The present invention is directed to the use of certain pyrido[1,2-a]-pyrazine derivatives, also described as bis-azabicyclic compounds, in the treatment of Parkinson's disease, attention deficit hyperactivity disorder ("ADHD") and microadenomas in mammals, including humans. It is also directed to the use of a dopamine-2 (D2) receptor agonist in combination with a serotonin-1A (5HT_{1A}) receptor agonist for the treatment for Parkinson's Disease. It is also directed to the use of an alpha-2 (α_2) adrenergic receptor ligand in combination with either a D2 receptor agonist or a 5HT_{1A} receptor agonist for the treatment of ADHD. It is also directed to the use of a D2 receptor agonist in combination with a 5HT_{1A} receptor agonist for the treatment of ADHD. It is also directed to the use of an alpha-2 (α_2) adrenergic receptor ligand in combination with both a D2 receptor agonist and a 5HT_{1A} receptor agonist for the treatment of ADHD.

Background of the Invention

Serotonin plays a role in several psychiatric disorders, including anxiety, Alzheimer's disease, depression, nausea and vomiting, eating disorders, and migraine. (See Rasmussen *et al.*, "Chapter 1. Recent Progress in Serotonin (5HT)_{1A} Receptor Modulators", in Annual Reports in Medicinal Chemistry, Section I, 30, pp. 1-9, 1995, Academic Press, Inc.; Antigas *et al.*, Trends Neurosci., 19 (9), 1996, pp. 378-383; and Wolf *et al.*, Drug Development Research, 40, 1997, pp. 17-34.) Serotonin also plays a role in both the positive and negative symptoms of schizophrenia. (See Sharma *et al.*, Psychiatric Annals, 26 (2), February, 1996, pp. 88-92.) Serotonin 1A receptor agonists have been shown to increase prefrontal cortex dopamine (DA) release. See Wedzony *et al.*, Eur. J. Pharmacol., 305: 73-78 (1996). Buspirone, a 5HT_{1A} receptor agonist, has been shown to be efficacious in treating a variety of symptoms associated with ADHD. Serotonin 1A receptor agonists have also been shown to reverse neuroleptic induced dystonia in nonhuman primates, a condition that mimics symptoms of human Parkinson's disease. See Casey, D. E., Neuropsychopharmacol., 10:370S (1994).

Symptoms associated with ADHD have been shown to be relieved by catecholamine releasing drugs such as methylphenidate, and by postsynaptic α_2 adrenergic receptor agonists such as clonidine. Also, presynaptic α_2 adrenergic receptor antagonists have been shown to increase norepinephrine (NE) release.


A number of 1-(2-pyrimidinyl)-4-[4-(cyclic-imido)butyl]piperidine derivatives have been disclosed as anxiolytic agents which are generally lacking sedative activity. Among these are buspirone, where the cyclic-imido group is 4,4-tetramethylene-piperidine-2,6-dione-1-yl (Wu *et al.*, United States Patents 3,717,634 and 3,907,801); Casten *et al.*, United States Patent

5 4,182,763); gepirone, where the group is 4,4-dimethylpiperidine-2,6-dion-1-yl (Temple, Jr., United States Patent 4,423,049); and ipsapirone, where the group is 1,1-dioxobenzo[d] isothiazol-3(2H)-on-yl (Dompert *et al.*, German patent publication 3,321,969-A1). See also Ishizumi *et al.*, United States Patents 4,507,303 and 4,543,55; Freed *et al.*, United States Patent 4,562,255; Stack *et al.*, United States Patent 4,732,983; New *et al.*, United States
 10 Patent 4,524,026; and Stack, United States Patent 4,788,290.

Compounds of the formula (I) below are disclosed in United States Patent 5,122,525 as useful for the treatment of anxiety and depression. The use of such compounds for the treatment of addiction is described in United States Patent 5,616,885.

Summary of the Invention

15 This invention relates to a method of treating a disorder selected from Parkinson's disease, ADHD, and microadenomas in a mammal, including a human, comprising administering to a mammal in need of such treatment an amount of a compound of the formula






20


or a pharmaceutically acceptable acid addition salt thereof, wherein

X is N or CH;

Y is



or



5 Z is




SCH₂, OCH₂, Y¹(CH₂)_n or Y¹(CH₂)_n substituted on carbon with up to 2 methyl groups;
n is 1 or 2; and

Y¹ is CH₂, NH or NCH₃;

10 that is effective in treating such disorder.

In the compounds of the formula (I), Y is preferably



A particularly preferred compound is that wherein Z is Y¹(CH₂)_n, Y¹ is CH₂, n is 1 and
X is N.

15 The compounds of formula I are D₂ receptor agonists and useful in the treatment of
Parkinson's disease. They also exhibit 5HT_{1A} receptor agonist activity.

The compounds of formula I also exhibit activity as α₂ adrenergic receptor antagonists
and are useful in treatment of ADHD. The compounds increase hippocampal NE release and
also increase prefrontal cortex DA release.

20 The compounds of the formula I that are basic can form acid addition salts with a variety
of organic and inorganic acids. The acids that can be used to prepare the pharmaceutically
acceptable acid addition salts of the aforementioned base compounds of the formula I are those
which form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable
anions, such as the hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate,
25 phosphate, acid phosphate, acetate, lactate, citrate, acid citrate, tartrate, bitartrate, succinate,
maleate, fumarate, gluconate, saccharate, benzoate, methanesulfonate, ethanesulfonate,
benzenesulfonate, p-toluenesulfonate and pamoate [i.e., 1,1'-methylene-bis-(2-hydroxy-3-
naphthoate)]salts.

30 The term "treating" as used herein, refers to reversing, alleviating, inhibiting the progress
of, or preventing the disorder or condition to which such term applies, or one or more symptoms
of such disorder or condition. The term "treatment", as used herein, refers to the act of treating,
as "treating" is defined immediately above.

5 This invention also relates to a method for treating Parkinson's Disease in a mammal, including a human, comprising administering to a mammal in need of such treatment a D2 receptor agonizing agent in combination with a 5HT_{1A} receptor agonizing agent, wherein the two foregoing active agents are present in amounts such that the combination of such agents is effective in treating Parkinson's Disease.

10 This invention also relates to a pharmaceutical composition for treating Parkinson's disease in a mammal, including a human, comprising: (a) a D2 receptor agonizing agent or a pharmaceutically acceptable salt thereof; (b) a 5HT_{1A} receptor agonizing agent or pharmaceutically acceptable salt thereof; and (c) a pharmaceutically acceptable carrier, wherein the foregoing two active agents are in the composition in amounts such that the combination of
15 such agents is effective in treating Parkinson's disease.

This invention also relates to a method of treating ADHD in a mammal, including a human, comprising administering to a mammal in need of such treatment an α_2 adrenergic receptor ligand, or pharmaceutically acceptable salt thereof, in combination with either a D2 receptor agonizing agent or a 5HT_{1A} receptor agonizing agent, or pharmaceutically acceptable
20 salt thereof, wherein the foregoing two active agents are present in amounts such that the combination of such active agents is effective in treating ADHD.

This invention also relates to a pharmaceutical composition for treating ADHD in a mammal, including a human, comprising: (a) an α_2 adrenergic receptor ligand or a pharmaceutically acceptable salt thereof; (b) a D2 receptor agonizing agent or a 5HT_{1A} receptor
25 agonizing agent, or a pharmaceutically acceptable salt thereof; and (c) a pharmaceutically acceptable carrier, wherein the two foregoing active agents are present in the composition in amounts such that the combination of such agents is effective in treating ADHD.

This invention also relates to a method of treating ADHD in a mammal, including a human, comprising administering to a mammal in need of such treatment a D2 receptor
30 agonizing agent, or pharmaceutically acceptable salt thereof, in combination with a 5HT_{1A} receptor agonizing agent, or pharmaceutically acceptable salt thereof, wherein the foregoing two active agents are present in amounts such that the combination of such active agents is effective in treating ADHD.

This invention also relates to a pharmaceutical composition for treating ADHD in a mammal, including a human, comprising: (a) a D2 receptor agonizing agent or a
35 pharmaceutically acceptable salt thereof; (b) a 5HT_{1A} receptor agonizing agent or a pharmaceutically acceptable salt thereof; and (c) a pharmaceutically acceptable carrier, wherein the two foregoing active agents are present in the composition in amounts such that the combination of such agents is effective in treating ADHD.

40 This invention also relates to a method of treating ADHD in a mammal, including a human, comprising administering to a mammal in need of such treatment an α_2 adrenergic

5 receptor ligand, or pharmaceutically acceptable salt thereof, in combination with a D2 receptor agonizing agent, or pharmaceutically acceptable salt thereof, and also a 5HT_{1A} receptor agonizing agent, or pharmaceutically acceptable salt thereof, wherein the three foregoing active agents are present in amounts such that the combination of such active agents is effective in treating ADHD.

10 This invention also relates to a pharmaceutical composition for treating ADHD in a mammal, including a human, comprising: (a) an α_2 adrenergic receptor ligand or a pharmaceutically acceptable salt thereof; (b) a D2 receptor agonizing agent or a pharmaceutically acceptable salt thereof; (c) a 5HT_{1A} receptor agonizing agent or a pharmaceutically acceptable salt thereof; and (d) a pharmaceutically acceptable carrier; wherein
15 the foregoing three active agents are present in the composition in amounts such that the combination of such agents is effective in treating ADHD.

Compounds of the formula I may contain chiral centers and therefore may exist in different enantiomeric and diastereomeric forms. The term "compounds of the formula I", as used herein, refers to all optical isomers and all other stereoisomers of compounds of the formula I, as
20 defined above, and to all racemic and other mixtures thereof, and to all pharmaceutical compositions and methods of treatment defined above that contain or employ such isomers or mixtures.

Formula I above includes compounds identical to those depicted but for the fact that one or more hydrogen or carbon atoms are replaced by isotopes thereof. Such compounds are
25 useful as research and diagnostic tools in metabolism pharmacokinetic studies and in binding assays. Specific applications in research include radioligand binding assays, autoradiography studies and *in vivo* binding studies.

Examples of D2 receptor agonizing agents that can be used in the methods of this invention include, but are not limited to, compounds of the formula I and the pharmaceutically
30 acceptable salts, pergolide, bromocriptane, ropinerol, and pramipexole.


Examples of 5HT_{1A} receptor agonizing agents that can be used in the methods of this invention include, but are not limited to: (a) compounds of the formula I and their pharmaceutically acceptable salts, (b) buspirone (U.S. Patents 3,717,638; 3,907,801 and
35 4,182,763); (c) gepirone (U.S. Patent 4,423,049); (c) ipsapirone (German patent publication 3,321,969-A1); and (d) and flexinoxan.

Examples of α_2 adrenergic receptor antagonizing agents that can be used in the methods of this invention include, but are not limited to, compounds of the formula I and their pharmaceutically acceptable salts, yohimbine and idazoxan.

Examples of more specific embodiments of this invention are the above methods of
40 treating ADHD and pharmaceutical compositions for the treatment of ADHD that employ an α_2


- 5 adrenergic receptor ligand wherein such ligand is a presynaptic α_2 adrenergic receptor antagonist or a post synaptic α_2 adrenergic receptor agonist.

A preferred method of this invention is a method of treating Parkinson's Disease in a mammal, including a human, comprising administering to a mammal in need of such treatment an amount of the following compound of the formula I shown below:



10

i.e., wherein X is nitrogen, wherein Y is a group of the formula



("sunipetron"), or pharmaceutically acceptable salt thereof, that is effective in treating Parkinson's Disease.

15

Another preferred embodiment of this invention is a method for treating ADHD in a mammal including a human, comprising administering to a mammal in need of such treatment an amount of sunipetron, or pharmaceutically acceptable salt thereof, that is effective in treating ADHD.

20

Another preferred method of this invention is a method for treating microadenomas in a mammal including a human, comprising administering to a mammal in need of such treatment an amount of sunipetron, or pharmaceutically acceptable salt thereof, that is effective in treating microadenomas.

Brief Description of the Drawings

Figure 1 shows parkinsonian scores obtained in monkeys treated with either 0.5 mg/kg or 1.0 mg/kg of a compound of formula I.

25

Detailed Description of the Invention

All patents, patent applications, and articles listed herein are hereby incorporated by reference in their entireties.

Compounds of the formula I and their pharmaceutically acceptable salts may be prepared as described in United States Patent 5,122,525; United States Patent 5,185,449,

30

- 5 United States Patent 5,455,350, United States Patent Application 08/470,377, which was filed on June 6, 1995, and World Patent Application PCT/IB97/00704, which designates the United States and was filed on June 16, 1997.

The compounds of the formula I are capable of forming a wide variety of different salts with various inorganic and organic acids. The acids that can be used to prepare the
10 pharmaceutically acceptable acid addition salts of the base compounds of this invention are those which form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, such as hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate or bisulfate, phosphate or acid phosphate, acetate, lactate, citrate or acid citrate, tartrate or bitartrate, succinate,
15 maleate, fumarate, gluconate, saccharate, benzoate, methanesulfonate and pamoate [i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)] salts. Although such salts must be pharmaceutically acceptable for administration to animals, it is often desirable in practice to initially isolate a compound of the formula I from the reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base compound by treatment with an alkaline reagent, and subsequently convert the free base to a pharmaceutically acceptable acid addition
20 salt. The acid addition salts of the base compounds of this invention are readily prepared by treating the base compound with a substantially equivalent amount of the chosen mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent such as methanol or ethanol. Upon careful evaporation of the solvent, the desired solid salt is obtained.

For use in treating Parkinson's disease, ADHD or microadenomas in a human
25 subject, a compound of the formula I, or a pharmaceutically-acceptable salt thereof, is administered in an amount of about 2-300 mg/day, in single or divided daily doses. In particular cases, dosages outside that range are prescribed at the discretion of the attending physician. The preferred route of administration is generally oral, but parenteral administration (e.g., intramuscular, intravenous, intradermal) will be preferred in special cases, e.g., where
30 oral absorption is impaired as by disease, or the patient is unable to swallow. In one embodiment, the compound of formula I is administered in an individual dosage of about 1.0 mg/kg at a frequency of about three times per day. A controlled-release formulation can be employed instead that is administered once per day. In another embodiment, the compound is administered in an amount required by the particular route of administration to achieve a
35 plasma concentration of between 100 and 500 ng/ml, preferably about 200 ng/ml, at between 2 and 3 hours following administration.

The compounds used in the present invention are generally administered in the form of pharmaceutical compositions comprising at least one of the compounds of the formula (I), or a salt thereof, together with a pharmaceutically acceptable vehicle or diluent. Such
40 compositions are generally formulated in a conventional manner utilizing solid or liquid vehicles or diluents as appropriate to the mode of desired administration: for oral administration, in the form of tablets, hard or soft gelatin capsules, suspension, granules,

- 5 powders and the like; and, for parenteral administration, in the form of injectable solutions or suspensions, and the like.

Binding of a compound to the D₂ receptor can be determined using the follow D₂ receptor binding assay.

- 10 LTK cells expressing the human D₂ long (D_{2L}) receptor are grown (T-175 flasks) in D-glucose containing minimal essential media (DMEM; Gibco) supplemented with 10% fetal bovine serum (FBS). The cells are dislodged with 5 mM EDTA in PBS and homogenized in 50 mM Tris HCl (pH 7.4) with 5mM MgSO₄, using a Brinkman Polytron at setting 6 for 20 sec. Membranes are recovered after multiple rounds of separation by centrifugation and resuspension in fresh ice-cold buffer. The tissue (~ 2 mg tissue, wet weight) is added to test
- 15 tubes containing incubation buffer (50 mM Tris HCl, 120 mM NaCl, 2 mM MgCl₂, 5 mM KCl, 5 mM CaCl₂, pH 7.2), various concentrations of test drug, and [³H]-spiperone (0.06 nM final concentration, Amersham, Arlington Heights IL). Non-specific binding is determined in the presence of 2 uM (+)-butaclamol. After 45 min at 30°C, incubations are terminated by rapid filtration through Whatman GF/B filters using a Brandel cell harvester. The membranes are
- 20 washed using 3 X 4 ml of ice-cold buffer and membrane-bound ligand is determined by liquid scintillation counting of the filters in Ready-Safe scintillation cocktail (for tritiated ligands). The K_d (0.06 nM) for the radioligand is determined previously by saturation analysis and used to calculate apparent K_i's by means of the Cheng-Prusoff equation.

- 25 The agonist or antagonist activity of a compound at the D₂ receptor can be determined using the following three assays.

(1) Human D₂ receptor modulation of cAMP formation in GH₄C₁ cells

- GH₄C₁ cells, derived from rat pituitary, expressing either the long or short forms of the human D₂ receptor, are grown to confluence in (HAM) F-10 Nutrient Mixture (Gibco) supplemented with 10% FBS and 2 mM l-glutamine and 10 U/ml penicillin-streptomycin in T-
- 30 175 flasks. The cells are dislodged with 5 mM ethylenediamine tetraacetic acid (EDTA) in phosphate buffered saline (PBS) and resuspended in PBS containing 5 mM MgCl₂, 30 mM hydroxyethylpiperazine-N-ethanesulfonic acid (HEPES), and 50 mM isobutyl methyl xanthine (IBMX). Cells (~200,000/tube) are exposed to 5 mM forskolin, 100 nM quinpirole or forskolin plus quinpirole plus antagonist for 11 minutes. In experiments with antagonists, cells are
- 35 exposed to the antagonists 11 minutes prior to quinpirole challenge. To judge agonist activity, the effect of a compound on forskolin-stimulated cAMP accumulation is tested in the absence of the agonist quinpirole. The reaction is terminated with the addition of 6N perchloric acid, and samples are neutralized with 5N potassium hydroxide and 2 M Tris buffer. Cyclic AMP levels are measured using a commercially available competitive binding kit (Amersham). IC₅₀
- 40 values are calculated by linear regression analysis of the concentration-response curves. Apparent K_i values are calculated using the equation: $K_i = IC_{50} / (1 + [agonist] / [agonist EC_{50}])$.

(2) Electrophysiology in Rat Brain Slices

5 Male Sprague-Dawley rats (200-250 gm, Charles River Laboratories, Wilmington, MA) are lightly anesthetized with halothane, decapitated and the brains quickly removed to ice-cold, oxygenated medium (95% O₂/5% CO₂; 124 mM NaCl, 2 mM KCl, 1.25 mM NaH₂PO₄, 26 mM NaHCO₃, 10 mM d-glucose, 2 mM MgSO₄·7H₂O and 2 mM CaCl₂; pH 7.4). The ventral tegmental area is blocked and glued using cyanoacrylate to the stage of a Lancer
10 Vibratome (Series 1000) filled with ice cold medium. Coronal slices (350 µ) are cut and placed in oxygenated medium (22°C) for 1 hour prior to recording. For recording, slices are placed on a nylon net in a recording chamber where they are completely submerged in continually flowing medium at 35°C (~1 ml/min). All drugs are applied by switching the perfusion medium to a solution containing the drug. Spontaneous extracellular action
15 potentials are recorded using 0.9% saline-filled glass pipettes (6-8 MW). Firing rates are plotted on-line in bins of 10 or 20 seconds and alterations in firing rate are calculated using average rates over 2 min epochs before and after drug application. Concentration-response curves are constructed and analyzed by linear regression.

(3) Microdialysis

20 I shaped concentric microdialysis probes are constructed out of dialysis fiber (molecular weight cut off of 18,000, 300 µm o.d., Hospal, The Netherlands) occluded at one end with epoxy resin and attached to fused silica microtubing. The probes, 9 mm long with 2 mm length of exposed dialysis membrane are implanted into the nucleus accumbens (AP 1.7, ML -1.2, DV -8.0) of male Sprague Dawley rats (300-350g) anesthetized with ketamine (75
25 mg/kg) and xylazine (10 mg/kg). Following surgery, rats are placed in perspex cages inside insulation boxes and the probe inlets connected via flexible PEEK tubing through a dual channel fluid swivel system to a CMA/100 microinfusion pump (CMA/Microdialysis, Acton, MA). The probe is perfused overnight with artificial cerebrospinal fluid (147 mM NaCl₂, 2.7 mM KCl, 1.3 mM CaCl₂, 1.0 mM MgCl₂ and 0.1 mM ascorbic acid) at 0.5 ml/min. The next
30 day, an experiment is started by increasing the flow to 1.5 ml/min and connecting the probe outlet with PEEK tubing to a 30 ml sample loop in a DECADE electrochemical detector (ANTEC, Leiden, The Netherlands). Microdialysis samples (30 µl) are collected on-line and automatically injected onto the column every 20 or 25 min. Analytes are separated at 35°C over a BDS Hypersil C₁₈ 3 m column (150 X3 mm) by reverse phase HPLC using a 75 mM
35 potassium phosphate mobile phase of pH 5.0, containing 0.8 mM octanesulfonate, 8% methanol, 3 mM triethylamine and delivered at a flow rate of 0.35 ml/min by an ESA 580 pump. Amperometric detection of dopamine is performed using a glassy carbon electrode of the DECADE detector set at 550 mV vs Ag/AgCl. Extracellular levels of dopamine are quantified by comparing peak heights with those of standards.

40 After obtaining a stable baseline (5-7 samples collected every 20 or 25 min) drugs are administered and release of DA monitored for 4 to 7 hours. Dialysate concentrations are expressed as a percentage of baseline. Dialysate concentrations of DA are not corrected for

5 recovery across the dialysis fiber. To determine whether each dose of drug has a significant effect on nucleus accumbens DA release multivariate analysis of variance with repeated measures over time is performed using SuperANova software (Abacus Concepts, Inc. Berkeley CA.)

10 The ability of a particular compound to exhibit an anti-Parkinsonian effect in nonhuman primates can be determined using the procedures described by Greenemyre *et al.*, *Ann. Neurol.*, 35:655-661, 1994, and Klockgether *et al.*, *Ann. Neurol.*, 30:717-723, 1991.

15 The agonist and antagonist activities of a particular compound of the invention at 5-HT_{1A} receptors can be determined using a single saturating concentration according to the following procedure. Male Hartley guinea pigs are decapitated and 5-HT_{1A} receptors are dissected out of the hippocampus. The individual tissues are homogenized in 5 mM HEPES buffer containing 1 mM EGTA (pH 7.5) using a hand-held glass-Teflon® homogenizer and centrifuged at 35,000 x g for 10 minutes at 4°C. The pellets are resuspended in 100 mM HEPES buffer containing 1 mM EGTA (pH 7.5) to a final protein concentration of 20 mg (hippocampus) or 5 mg (substantia nigra) of protein per tube. The following agents are added
20 so that the reaction mix in each tube contained 2.0 mM MgCl₂, 0.5 mM ATP, 1.0 mM cAMP, 0.5 mM IBMX, 10 mM phosphocreatine, 0.31 mg/mL creatine phosphokinase, 100 mM GTP and 0.5-1 microcuries of [³²P]-ATP (30 Ci/mmol: NEG-003 - New England Nuclear). Incubation is initiated by the addition of tissue to siliconized microfuge tubes (in triplicate) at 30°C for 15 minutes. Each tube receives 20 mL tissue, 10 mL drug or buffer (at 10X final concentration), 10mL 32 nM agonist or buffer (at 10X final concentration), 20mL forskolin (3
25 mM final concentration) and 40 mL of the preceding reaction mix. Incubation is terminated by the addition of 100 mL 2% SDS, 1.3 mM cAMP, 45 mM ATP solution containing 40,000 dpm [³H]-cAMP (30 Ci/mmol: NET-275 - New England Nuclear) to monitor the recovery of cAMP from the columns. The separation of [³²P]-ATP and [³²P]-cAMP is accomplished using the
30 method of Salomon *et al.*, *Analytical Biochemistry*, 1974, 58, 541-548. Radioactivity is quantified by liquid scintillation counting. Maximal inhibition is defined by 10 mM (R)-8-OH-DPAT for 5-HT_{1A} receptors. Percent inhibition by the test compound is then calculated in relation to the inhibitory effect of (R)-8-OH-DPAT. The reversal of agonist induced inhibition of forskolin-stimulated adenylate cyclase activity is calculated in relation to the 32 nM agonist
35 effect.

This invention relates both to methods of treating Parkinson's disease and ADHD in which the two or three active agents employed are administered together, as part of the same pharmaceutical composition, as well as to methods in which these active agents are administered separately as part of an appropriate dose regimen designed to obtain the benefits
40 of the combination therapy. The appropriate dose regimen, the amount of each dose administered, and specific intervals between doses of each active agent will depend upon the subject being treated, how well tolerated the drug is and the severity of the condition. Generally,

5 in carrying out the above combination methods of this invention, the 5HT_{1A} receptor agonist will be administered in an amount ranging from about 5 - 90 mg per day, in single or divided doses, the α_2 adrenergic receptor ligand will be administered in an amount ranging from about 1.0 - 100 mg per day, in single or divided doses, in the case of an α_2 adrenergic receptor antagonist, and
10 an α_2 adrenergic receptor agonist, in single or divided doses, and the D2 receptor agonist will be administered in an amount ranging from about 0.5 mg - 25 mg per day, in single or divided doses. (For FDA approved drugs that are used in the combination methods of this invention, physicians will be guided by the dosage ranges for such drugs that are specified in the Physician's Desk Reference). Variations may nevertheless occur depending upon the species of
15 animal being treated and its individual response to said medicament, as well as on the type of pharmaceutical formulation chosen and the time period and interval at which such administration is carried out. In some instances, dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effect, provided that such larger doses are first divided into several small doses
20 for administration throughout the day.

The combination methods of this invention include methods wherein the desired combined activities are present in one compound or pharmaceutically acceptable salt. The pharmaceutical compositions of this invention that exhibit more than one pharmaceutical activity (e.g., 5HT_{1A} agonism and D2 agonism) include those pharmaceutical compositions wherein all
25 the desired pharmaceutical activities are present in one compound or pharmaceutically acceptable salt.

The D2 receptor agonists, the 5HT_{1A} receptor agonists, and the α_2 receptor antagonists that are employed in the pharmaceutical compositions and methods of this invention are hereinafter also referred to as "therapeutic agents". The therapeutic agents can be administered
30 via either the oral or parenteral route. Compositions containing both a D2 receptor agonist and a 5HT_{1A} receptor agonist, or all three of a D2 receptor agonist, a 5HT_{1A} receptor agonist and an α_2 adrenergic receptor antagonist, will generally be administered orally or parenterally daily, in single or divided doses, so that the total amount of each active agent administered falls within the above guidelines.

35 The therapeutic agents may be administered alone or in combination with pharmaceutically acceptable carriers or diluents by either of the routes previously indicated, and such administration may be carried out in single or multiple doses. More particularly, the therapeutic agents of this invention can be administered in a wide variety of different dosage forms, i.e., they may be combined with various pharmaceutically acceptable inert carriers in the
40 form of tablets, capsules, lozenges, troches, hard candies, suppositories, aqueous suspensions, injectable solutions, elixirs, syrups, and the like. Such carriers include solid diluents or fillers,

5 sterile aqueous media and various non-toxic organic solvents, etc. Moreover, oral pharmaceutical compositions can be suitably sweetened and/or flavored. In general, the therapeutic compounds of this invention, when administered separately (*i.e.*, not in the same pharmaceutical composition) are present in such dosage forms at concentration levels ranging from about 5.0% to about 70% by weight.

10 For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dicalcium phosphate and glycine may be employed along with various disintegrants such as starch (and preferably corn, potato or tapioca starch), alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as
15 magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral
20 administration, the active ingredient may be combined with various sweetening or flavoring agents, coloring matter or dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

For parenteral administration, solutions of a therapeutic agent in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be
25 suitably buffered if necessary and the liquid diluent first rendered isotonic. These aqueous solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intraarticular, intramuscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well known to those skilled in the art.

30 The following example is intended only to illustrate the invention, and not to be interpreted as limiting its scope.

Example

The behavioral effects of administration of sunipetron on parkinsonian score in MPTP- (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine) treated monkeys was determined. The
35 methods employed are described in Greenemyre et al., *supra* and Klockgether et al., *supra*.

Briefly, parkinsonian monkeys were administered 0.5 mg/kg or 1.0 mg/kg of sunipetron and the severity of parkinsonian symptoms was rated at each of 4 time points after drug administration as compared with administration of a control solution ("Vehicle").


Parkinsonian scores obtained from the study are shown in Figure 1. Combined with
40 the estimates shown for plasma levels of the compound following administration, the data demonstrate that the maximum reduction in parkinsonian score was attained at plasma levels

- 5 of approximately 200 ng/ml. The minimum effective plasma level was approximately 100 ng/ml.

CLAIMS

5

1. A method of treating a disorder selected from Parkinson's Disease, ADHD, and microadenomas in a mammal, comprising administering to a mammal in need of such treatment an amount of a compound of the formula






10


or a pharmaceutically acceptable acid addition salts thereof, wherein

X is N or CH;


Y is



or



Z is



15


SCH_2 , OCH_2 , $\text{Y}^1(\text{CH}_2)_n$ or $\text{Y}^1(\text{CH}_2)_n$ substituted on carbon with up to 2 methyl groups;

n is 1 or 2; and

Y^1 is CH_2 , NH or NCH_3 ;

that is effective in treating such disorder.

- 5 2. A method according to claim 1 wherein the compound of formula (I) or pharmaceutically acceptable salt that is employed is one wherein Y is



Z is $Y^1(CH_2)_n$, Y^1 is CH_2 , n is 1 and X is N.

- 10 3. A method for treating Parkinson's disease in a mammal, comprising administering to a mammal in need of such treatment a D2 receptor agonizing agent, or a pharmaceutically acceptable salt thereof, in combination with a $5HT_{1A}$ receptor agonizing agent, or a pharmaceutically acceptable salt thereof, wherein the two foregoing active agents are present in amounts such that the combination of such agents is effective in treating Parkinson's disease.

- 15 4. A pharmaceutical composition for treating Parkinson's disease in a mammal, comprising: (a) a D2 receptor agonizing agent, or a pharmaceutically acceptable salt thereof; (b) a $5HT_{1A}$ receptor agonizing agent, or a pharmaceutically acceptable salt thereof; and (c) a pharmaceutically acceptable carrier; wherein the foregoing two active agents are present in the composition in amounts such that the combination of such agents is effective in treating Parkinson's disease.

- 20 5. A method of treating ADHD in a mammal, comprising administering to a mammal in need of such treatment an α_2 adrenergic receptor ligand, or a pharmaceutically acceptable salt thereof, in combination with a D2 receptor agonizing agent, or a pharmaceutically acceptable salt thereof, and also in combination with a $5HT_{1A}$ receptor agonizing agent, or a pharmaceutically acceptable salt thereof, wherein the three foregoing active agents are present in amounts such that the combination of such active agents is effective in treating ADHD.

- 30 6. A pharmaceutical composition for treating ADHD in a mammal, comprising: (a) an α_2 adrenergic receptor ligand, or pharmaceutically acceptable salt thereof; (b) a D2 receptor agonizing agent, or a pharmaceutically acceptable salt thereof; (c) a $5HT_{1A}$ receptor agonizing agent, or pharmaceutically acceptable salt thereof; and (d) a pharmaceutically acceptable carrier; wherein the foregoing three active agents are present in the composition in amounts such that the combination of such active agents is effective in treating ADHD.

- 35 7. A method of treating ADHD in a mammal, comprising administering to a mammal in need of such treatment an α_2 adrenergic receptor ligand, or a pharmaceutically acceptable salt thereof, in combination with a D2 receptor agonizing agent or a $5HT_{1A}$ receptor agonizing agent, or a pharmaceutically acceptable salt thereof, wherein the foregoing two active

5 agents are present in amounts such that the combination of such active agents is effective in treating ADHD.

8. A pharmaceutical composition for treating ADHD in a mammal, comprising: (a) an α_2 adrenergic receptor ligand or a pharmaceutically acceptable salt thereof; (b) a D2 receptor agonizing agent or a 5HT_{1A} receptor agonizing agent, or a pharmaceutically acceptable salt thereof; and (c) a pharmaceutically acceptable carrier; wherein the two foregoing active agents
10 are in the composition in amounts such that the combination of such agents in the composition is effective in treating ADHD.

9. A method of treating ADHD in a mammal, comprising administering to a mammal in need of such treatment a D2 receptor agonizing agent, or pharmaceutically acceptable salt thereof, in combination with a 5HT_{1A} receptor agonizing agent, or pharmaceutically acceptable salt thereof, wherein the foregoing two active agents are present in
15 amounts such that the combination of such active agents is effective in treating ADHD.

10. A pharmaceutical composition for treating ADHD in a mammal, comprising: (a) a D2 receptor agonizing agent or a pharmaceutically acceptable salt thereof; (b) a 5HT_{1A} receptor agonizing agent or a pharmaceutically acceptable salt thereof; and (c) a pharmaceutically acceptable carrier; wherein the foregoing two active agents are present in the composition in amounts such that the combination of such agents in the composition is effective
20 in treating ADHD.

11. A method according to claim 1 of treating Parkinson's disease in a mammal, comprising administering to a mammal in need of such treatment an amount of sunipetron, or a pharmaceutically acceptable salt thereof, that is effective in treating Parkinson's Disease.
25

12. A method according to claim 1 of treating ADHD in a mammal, comprising administering to a mammal in need of such treatment an amount of sunipetron, or a pharmaceutically acceptable salt thereof, that is effective in treating ADHD.

13. A method according to claim 1 of treating microadenomas in a mammal, comprising administering to a mammal in need of such treatment an amount of sunipetron, or a pharmaceutically acceptable salt thereof, that is effective in treating microadenomas.
30


14. A method according to claim 5, wherein the α_2 adrenergic receptor ligand that is employed is a presynaptic α_2 adrenergic receptor antagonist or a postsynaptic α_2 adrenergic receptor agonist.
35

15. A method according to claim 7, wherein the α_2 adrenergic receptor ligand that is employed is a presynaptic α_2 adrenergic receptor antagonist or a postsynaptic α_2 adrenergic receptor agonist.

16. A pharmaceutical composition according to claim 6, wherein the α_2 adrenergic receptor ligand that is employed is a presynaptic α_2 adrenergic receptor antagonist or a postsynaptic α_2 adrenergic receptor agonist.
40

5 17. A pharmaceutical composition according to claim 8, wherein the α_2 adrenergic receptor ligand that is employed is a presynaptic α_2 adrenergic receptor antagonist or a postsynaptic α_2 adrenergic receptor agonist.




18. A pharmaceutical composition for treating a disorder selected from Parkinson's Disease, ADHD, and microadenomas in a mammal, comprising an amount of a compound of
10 the formula




or a pharmaceutically acceptable acid addition salt thereof, wherein

X is N or CH;

Y is





or




15

Z is




SCH₂, OCH₂, Y¹(CH₂)_n or Y¹(CH₂)_n substituted on carbon with up to 2 methyl groups;
n is 1 or 2; and

- 5 Y^1 is CH_2 , NH or NCH_3 ;
that is effective in treating such disorder, and a pharmaceutically acceptable carrier.
19. A pharmaceutical composition according to claim 18 wherein the compound of formula (I) or pharmaceutically acceptable salt that is employed is one wherein, Y is



- 10 Z is $Y^1(CH_2)_n$, Y^1 is CH_2 , n is 1 and X is N.

FIG. 1



INTERNATIONAL SEARCH REPORT

Internat'l Application No
PCT/IB 99/01503

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 A61K31/505 A61P25/16		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 616 585 A (BRIGHT GENE M) 1 April 1997 (1997-04-01) cited in the application abstract column 1, line 50 -column 2, line 59 column 5, line 44 -column 6, line 67 claims 1-18 <div style="text-align: center; margin-top: 20px;"> --- -/-- </div>	18,19
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex. </div>		
* Special categories of cited documents : <div style="display: flex;"> <div style="flex: 1;"> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="flex: 1;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*Z* document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-size: 1.2em;">18 November 1999</div>	Date of mailing of the international search report <div style="text-align: center; font-size: 1.2em;">21.01.00</div>	
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer <div style="text-align: center; font-size: 1.2em;">Taylor, G</div>	

INTERNATIONAL SEARCH REPORT

Intern. Patent Application No

PCT/IB 99/01503

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	JACKSON D.M. ET AL.: "Time course of bromocriptine induced excitation in the rat: behavioural and biochemical studies." NAUNYN-SCHMIEDEBERGS ARCHIVES OF PHARMACOLOGY, vol. 351, no. 2, 1995, pages 146-155, XP002122877 abstract introduction table 1 page 154, column 1, last paragraph ---	1-3,11
Y	WO 95 28933 A (BOOTS CO PLC ;EESTEVE LABOR DR (ES); HEAL DAVID JOHN (GB); NEEDHAM) 2 November 1995 (1995-11-02) abstract page 1, line 2 -page 3, line 20 page 10, line 25 - line 31 claims 1-3 ---	1-3,11
Y	PRAKASH, C. ET AL.: "Metabolism and excretion of a new anxiolytic drug candidate, CP-93,393, in healthy male volunteers" DRUG METABOLISM AND DISPOSITION, vol. 26, no. 5, 1998, pages 448-456, XP002122878 abstract figure 1 ---	1-3,11
A	US 5 281 607 A (BING GUOYING ET AL) 25 January 1994 (1994-01-25) abstract claims 1-4 -----	1-3,11, 18,19

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB 99/01503

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 3
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
See further information sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1 (in part), 2 (in part), 3, 11, 18 (in part), 19 (in part)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/IB 99/01503

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1 (in part), 2 (in part), 3, 11, 18 (in part),
19 (in part)

Methods and compositions for the treatment of Parkinson's disease

2. Claims: 1 (in part), 2 (in part), 5-10, 12, 14-17,
18 (in part), 19 (in part)

Methods and compositions for the treatment of attention deficit hyperactivity disorder (ADHD).

3. Claims: 1 (in part), 2 (in part), 13, 18 (in part),
19 (in part)

Methods and compositions for the treatment of microadenomas.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/IB 99/01503

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Present claim 3 relates to a method for the treatment involving administration of a D2 receptor agonising agent in combination with a 5HT1A receptor agonising agent. It is not clear from the wording of the claim whether this is meant to entail the use of two separate agents, or whether one agent fulfilling both these requirements also falls within the scope. Bromocriptine is a known agonist of both D2 and 5HT1A receptors (see Naunyn-Schmiedeberg's Archives of Pharmacology 1995, 351(2), 146); it is also known for the treatment of Parkinson's disease. As noted on Form PCT/ISA/206, a further lack of unity could therefore arise in this case.

However, since the gist of the application is the use of compounds of Formula (I) (cf. claim 1), and support/disclosure within the meaning of Art. 5 and 6 PCT is only provided for compounds of Formula (I) (the only example is that of sunepitron), the search for claim 3 has been restricted to methods for the treatment of Parkinson's disease involving the administration of compounds of Formula (I).

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 99/01503

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5616585 A	01-04-1997	WO 9200075 A	09-01-1992
		AU 651977 B	11-08-1994
		DE 69106516 D	16-02-1995
		DE 69106516 T	11-05-1995
		EP 0536173 A	14-04-1993
		AT 116545 T	15-01-1995
		AU 7909491 A	23-01-1992
		CA 2086335 A,C	30-12-1991
		DK 536173 T	01-05-1995
		ES 2066447 T	01-03-1995
		IE 64467 B	09-08-1995
		IL 98588 A	29-06-1995
		NZ 238750 A	29-09-1999
		WO 9200074 A	09-01-1992
WO 9528933 A	02-11-1995	AU 2137595 A	16-11-1995
		ES 2116880 A	16-07-1998
		ZA 9503237 A	28-12-1995
US 5281607 A	25-01-1994	AU 5354394 A	09-05-1994
		WO 9408579 A	28-04-1994